

Flavourful hadronic physics

B. El-Bennich^a, M. A. Ivanov^b and C. D. Roberts^{ac}

^aPhysics Division, Argonne National Laboratory, Argonne, Illinois 60439, USA

^bBogoliubov Laboratory of Theoretical Physics, JINR, 141980 Dubna, Russia

^cDepartment of Physics, Peking University, Beijing 100871, China

We review theoretical approaches to form factors that arise in heavy-meson decays and are hadronic expressions of non-perturbative QCD. After motivating their origin in QCD factorisation, we retrace their evolution from quark-model calculations to non-perturbative QCD techniques with an emphasis on formulations of truncated heavy-light amplitudes based upon Dyson-Schwinger equations. We compare model predictions exemplarily for the $F^{B \rightarrow \pi}(q^2)$ transition form factor and discuss new results for the $g_{D^* D \pi}$ coupling in the hadronic D^* decay.

1. Flavour physics and strong phases

The last two decades have witnessed important advances in flavour physics and in particular heavy-meson decays. From the first observation of a B meson by the CLEO Collaboration in 1981 at the Cornell Electron Storage Ring [1] (and their ongoing D -meson research program) to the dedicated B -physics facilities at SLAC in California and KEK in Japan, much progress has been made. Of course, while B physics is the main focus of the Collaborations Belle at KEK and BaBar at SLAC, and of the CDF experiment at Fermilab, considerable efforts have also been devoted to studies of D -meson decays, charmonium and τ physics.

Naturally, the driving force is to confirm the electroweak sector of the Standard Model which has established itself as the foremost paradigm to describe CP violation; in other words, the main task experimentalists strive for is the precise area of the Cabibbo-Kobayashi-Maskawa (CKM) triangle and the weak CP violating phase codified within its angles. In order to measure the size and exact form of this triangle, the angles are derived from branching fractions in a nowadays large variety of decay channels. We focus on non-leptonic decays, $B \rightarrow M_1 M_2$, but our discussion also applies to the cleaner semi-leptonic $B \rightarrow M \ell \bar{\nu}_\ell$ case.

From a theoretical point of view, heavy mesons can be used to test simultaneously all the mani-

festations of the Standard Model, namely the interplay between electroweak and strong interactions. They also provide an excellent playground to examine non-perturbative QCD effects already much studied in hadronic physics. It is noteworthy to remind that no CP -violating amplitude can be generated without strong phases. Suppose a heavy particle H decays into a mesonic final state $M = M_1 M_2 \dots$, $H \rightarrow M$, and that the Standard Model lagrangian contributes two terms (two Feynman diagrams) to this process. Then, the decay amplitude and its corresponding CP conjugate, written most generally, are

$$\mathcal{A}(H \rightarrow M) = \lambda_1 A_1 e^{i\varphi_1} + \lambda_2 A_2 e^{i\varphi_2} \quad (1)$$

$$\bar{\mathcal{A}}(\bar{H} \rightarrow \bar{M}) = \lambda_1^* A_1 e^{i\varphi_1} + \lambda_2^* A_2 e^{i\varphi_2} . \quad (2)$$

The weak coupling λ_i is a combination of possibly complex CKM matrix elements and $A e^{i\varphi}$ denotes the strong (hadronic) parts of the transition amplitude, where we emphasise that they too can have both a real part, or magnitude, and a phase, or absorptive part, due to multiple rescattering of the final-state quarks and mesons. These CP -related intermediate states must contribute the same absorptive part to the two decays, therefore the strong phases φ_i are the same in Eqs. (1) and (2). Taking the difference of the absolute squares, known as *direct* CP violation,

$$|\mathcal{A}|^2 - |\bar{\mathcal{A}}|^2 = 2A_1 A_2 \operatorname{Im}(\lambda_1 \lambda_2^*) \sin(\varphi_1 - \varphi_2) , \quad (3)$$

one sees that **no** such violation, $|\bar{\mathcal{A}}/\mathcal{A}| \neq 1$,

can occur if the weak couplings contain only real phases or the strong phases are the same. Hence, in order to extract the weak CKM phases with precision from the decay amplitudes, it is crucial to evaluate the QCD contributions *reliably*.

2. QCD factorisation

As simple as these mesons appear to be—a bound colourless heavy-light $\bar{q}q$ pair—the difference in quark masses and the energetic light mesons produced in their decays lead to a disparate array of energy scales. A central aspect of heavy-meson phenomenology are factorisation theorems which allow for a disentanglement of short-distance or *hard* physics, which encompasses electroweak interactions and perturbative QCD, from long-distance or *soft* physics, governed by non-perturbative hadronic effects. In the following, we illustrate the factorisation with non-leptonic decays of a heavy meson H .

In the hamiltonian formulation of heavy-quark effective theory (HQET) [2], in which amplitudes are expanded in powers of Λ_{QCD}/m_h and the heavy quark is a static particle in the limit $m_h \rightarrow \infty$, the $H \rightarrow M_1 M_2$ decay amplitude is given by

$$\mathcal{A} = \frac{G_F}{\sqrt{2}} \sum_p \lambda_p \sum_i C_i(\zeta) \langle M_1 M_2 | O_i | H \rangle(\zeta), \quad (4)$$

where $\lambda_p = V_{pb} V_{pk}^*$ ($p = u, c; k = d, s$) are products of CKM matrix elements and G_F is the Fermi coupling constant. The dimension-six four-quark operators O_i result from integrating out the weak gauge bosons $W^\pm$ in the operator product expansion and the Wilson coefficients $C_i(\zeta)$ encode perturbative QCD effects above the renormalisation point ζ .

In what is called “naive” factorisation, the hadronic matrix element $\langle M_1 M_2 | O_i | H \rangle$ is approximated by the product of two bilinear currents, $\langle M_1 | \bar{q} \gamma^\mu (1 - \gamma_5) b | B \rangle \otimes \langle M_2 | \bar{q}' \gamma_\mu (1 - \gamma_5) q | 0 \rangle + (M_1 \leftrightarrow M_2)$, where colour indices have been omitted. This factorisation simply expresses the matrix element of a local four-quark operator as a product of a decay constant and a transition form factor. However, as has long been known, the saturation by vacuum insertion fails in the case of most D decay modes and is largely in-

sufficient to reproduce the experimentally more precise data on $B \rightarrow M_1 M_2$ branching fractions. In fact, any hard final-state gluon interaction has been neglected and soft-gluon exchange is at best incorporated into an effective colour parameter or form factors. Moreover, the renormalisation scale and scheme dependence of $C_i(\zeta)$ are not cancelled by those of the matrix element $\langle M_1 M_2 | O_i | H \rangle(\zeta)$.

A major improvement over this simple factorisation *Ansatz* comes from the systematic reorganisation of weak and QCD interactions in the HQET. Three distinctive approaches have emerged in recent years: QCD factorisation (QCDF) [3], perturbative QCD (pQCD) [4] and soft-collinear effective theory (SCET) [5]. We here focus on the form factors that emerge in these factorisation approaches and solely remark that QCD corrections beyond naive factorisation entail, in the limit $m_h \gg \Lambda_{\text{QCD}}$, the $B \rightarrow M_1 M_2$ decay amplitude can be schematically written as

$$\begin{aligned} \langle M_1 M_2 | O_i | B \rangle &= \langle M_1 | j_1 | B \rangle \langle M_2 | j_2 | 0 \rangle \\ &\times \left[1 + \sum_n r_n \alpha_s^n + \mathcal{O}(\Lambda_{\text{QCD}}/m_b) \right], \quad (5) \end{aligned}$$

where j_1 and j_2 are the bilinear currents. This has been shown explicitly to leading order in α_s [3,6] and including the one-loop correction (α_s^2) to the tree-diagram scattering between the emitted meson and the one containing the spectator quark [7].

The factorisation theorem derived using SCET agrees with QCDF if perturbation theory is applied at the hard m_b^2 and hard-collinear $m_b \Lambda$ scales, with Λ typically of the order of 100 MeV. It is evident from Eq. (5) that higher orders in α_s break the factorisation but these corrections can be systematically supplemented; the analogy with perturbative factorisation for exclusive processes in QCD at large-momentum transfer is not accidental [8]. Further contributions that break the factorisation, formally suppressed in Λ_{QCD}/m_b yet not irrelevant, are weak annihilation decay amplitudes and final-state interactions between daughter hadrons [9]. Neglecting power corrections in α_s and taking the limit $m_b \rightarrow \infty$, the naive factorisation is recovered.

3. Separating scales: the softer the harder

While factorisation theorems elaborated with SCET provide the means to systematically integrate out energy scales in the perturbative domain, yielding approximations valid in the heavy-quark limit for a given decay in terms of products of hard and soft matrix elements, a reliable evaluation of the latter is notoriously difficult. In fact, it is the soft physics of the bound states that renders the task hard, as it implies non-perturbative QCD contributions. Full *ab initio* calculations are currently out of reach and for the time being one is left with modelling the heavy-to-light amplitudes with as much input from non-perturbative QCD as possible. Just how much soft physics is included depends on the theoretical *Ansatz* and techniques employed.

A variety of theoretical approaches have been applied to this problem, recent amongst which are analyses using light-front and relativistic quark models, light-cone sum rules (LCSR) and lattice-QCD simulations. In Section 4 we briefly summarise these approaches while Section 5 deals in more detail with studies of heavy-to-light form factors within the framework of the Dyson-Schwinger equation (DSE). We refer to a recent review [10] for a summary of transition form factor data from lattice-regularised QCD and just note that current results are obtained at large squared momentum transfer, *i.e.*, $q^2 \simeq 16 \text{ GeV}^2$ for $B \rightarrow \pi$ transitions. Hence, values at low q^2 must necessarily be extrapolated by means of a (pole-dominance) parametrisation [11,23].

4. Hadronic transition form factors

Quark models: relativistic quark models [12,13,14,15,16,17,18] have in common that their only degrees of freedom are constituent quarks whose masses are parameters of the hamiltonian. The hadronisation of the two valence quarks is described by vertex wave functions or phenomenological Bethe-Salpeter amplitudes (BSA). The approaches in [13,14,15] represent heavy-to-light transition amplitudes by triangle diagrams, a 3-point function between two meson BSA and the weak coupling, which yields the transition amplitude $\langle M(p_2) | \bar{q} \Gamma_I h | H(p_1) \rangle$ and reads generally,

$$\begin{aligned} \mathcal{A}(p_1, p_2) = & \text{tr}_{\text{CD}} \int \frac{d^4 k}{(2\pi)^4} \bar{\Gamma}_M^{(\mu)}(k; -p_2) S_q(k + p_2) \\ & \times \Gamma_I(p_1, p_2) S_Q(k + p_1) \Gamma_H(k; p_1) S_{q'}(k), \quad (6) \end{aligned}$$

where $S(k)$ are quark propagators, $Q = c, b$; $q = q' = u, d, s$; $M = S, P, V, A$ and the index μ indicates a possible vector structure in the final-state BSA. Γ_I is the interaction vertex whose Lorentz structure depends on the operator O_i in the HQET and Γ_H is the heavy meson BSA. The trace is over Dirac and colour indices.

An analogous approach represents the amplitude in Eq. (6) by relativistic double-dispersion integrals over the initial- and final-mass variables p_1^2 and p_2^2 , where the integration kernel arises from the double discontinuity of the triangle diagram (putting internal quark propagators on-shell via the Landau-Cutkosky rule). The meson-vertex functions are given by one-covariant BSA [16,17]. Other quark models [12] represent $B \rightarrow M$ form factors by overlap integrals of meson wave functions, obtained from confining potential models, and appropriate weak interaction vertices. Similar quark model calculations were performed on the light cone [18].

All these approaches model soft contributions with vertex functions, while the propagation of the constituent quark, $S(k) = (\not{k} - m_q)^{-1}$, is scale independent and does *not* describe confinement and dynamical chiral symmetry breaking (DCSB). As noted in Refs. [17,19,20], this can lead to considerable model dependence at larger momentum transfer.

Light-cone sum rules: In a LCSR the operator-product expansion of a given correlation function is combined with hadronic dispersion relations. The quark-hadron duality is invoked: the correlator function is calculated twice, as a hadronic object and with subhadronic degrees of freedom. After separation in HQET of the heavy meson's static part, $P_H = p + q = m_h v_h + k$, where v_h is the four-velocity and k is the residual momentum, and likewise redefinition of the momentum transfer $q = m_h v_h + \tilde{q} \Rightarrow p + \tilde{q} = k$, one obtains the heavy-limit correlation function,

$$\Pi^H(p, q) = \tilde{\Pi}^{H_v}(p, \tilde{q}) + \mathcal{O}(1/m_h); \quad (7)$$

q^2 [GeV ²]	Ref. [12]	Ref. [16]	Ref. [18]	Ref. [20]	Ref. [22]	Ref. [23]	Ref. [21]
0	0.217	0.29	0.247	0.24	0.25 ± 0.05	0.26 ± 0.03	$0.27 \pm 0.02 \pm 0.07$
10	0.41	0.54	0.49	0.53	0.54	0.51	–
15	0.67	0.84	0.97	0.91	0.90	0.81	–
20	1.40	1.56	> 10	1.75	1.83	1.58	–

Table 1

Numerical comparison for the transition form factor, $F_+^{B \rightarrow \pi}(q^2)$; the q^2 values of Ref. [20] are calculated, whereas for Refs. [12,16,18,22,23] the value $F_+^{B \rightarrow \pi}(0)$ and the corresponding extrapolation in these references are employed.

$$\tilde{\Pi}^{H_v}(p, \bar{q}) = i \int d^4x e^{ipx} \langle 0 | T[J_M(x) J_{h_v}(0)] | H_v \rangle,$$

whereas a hadronic correlator can be written,

$$\tilde{\Pi}^{\text{had.}}(p, q) = \frac{\langle 0 | J_M | M(p) \rangle \langle M(p) | J_h | H(P_H) \rangle}{m_M^2 - p^2}, \quad (8)$$

where $J_h(0)$ and $J_{h_v}(0)$ are heavy-light currents and $J_M(x)$ the interpolating current for a pseudoscalar or vector meson.

In Eq. (8) only the light-meson contribution is represented but higher and continuum states can also be taken into account. In Eqs. (7) and (8), the usual role of the correlation functions has been reversed [21,22]: the correlation function is taken between the vacuum and the on-shell B -state vector using its light-cone distribution amplitude (DA) expansion and the pion is interpolated with the light-quark (axialvector) current $J_M(x)$. The B -meson DAs are universal non-perturbative objects introduced within HQET.

In Ref. [23], however, the correlation function is taken between the vacuum and the light-meson state, whereas the B meson is interpolated by the heavy-light quark current $J_h(0)$. As a result, the long-distance dynamics in the correlation function is described by a set of light-meson (π, K, ρ, K^*) DA. In the last step, a Borel transformation is applied to both, Eqs. (7) and (8), from which one derives a transition form factor expressed as a sum rule. The transformation introduces a scale via the Borel parameter which, in turn, is fixed with sum rules for light-meson decay constants.

Besides a systematic uncertainty owing to the duality assumption, the main incertitude lies within the DAs which encode the *relevant* non-perturbative effects. Only their asymptotic form is known exactly from perturbative QCD. As of

yet, the first two Gegenbauer moments of the DA for various light pseudoscalar and vector mesons have been obtained from QCD sum rules with very large errors, though the first moment is consistent with lattice calculations [24]. Moreover, the transition form factors must be extrapolated to space-like momenta.

For purpose of comparison, we list $B \rightarrow \pi$ transitions form factors, $F_+(q^2)$, for various models in Table 1. As observed therein, there is a 30% variation within the quark models [12,16,18] at $q^2 = 0$ which increases at larger q^2 values. The LCSR predictions [22,23] agree at $q^2 = 0$ but their respective slopes for $q^2 > 0$ vary by 12%. The form factors obtained with the DSE model Ref. [20] are calculated on the entire physical momentum domain and the chiral limit is directly accessible.

5. Flavourful Dyson-Schwinger equations

The elements entering the amplitude in Eq. (6) can be motivated by the solutions of DSEs applied to QCD. A general review of the DSEs can be found in Refs. [25,26] and their applications to heavy-light transition form factors have been investigated in [20,27,28].

Dressed quark propagator: The mesons are bound states of a quark and antiquark pair, where for a given quark flavour their dressing is described by the DSE (in Euclidean metric),

$$S^{-1}(p) = Z_2(i\gamma \cdot p + m^{\text{bm}}) + \Sigma(p^2), \quad (9)$$

with the dressed quark self energy,

$$\Sigma(p^2) = Z_1 g^2 \int_k^\Lambda D^{\mu\nu}(p-k) \frac{\lambda^a}{2} \gamma_\mu S(k) \Gamma_\nu^a(k, p), \quad (10)$$

where $\int_k^\Lambda$ represents a Poincaré invariant regularisation of the integral with the regularisation

mass scale Λ . The current quark bare mass m^{bm} receives corrections from the self energy $\Sigma(p^2)$ in which the integral is over the dressed gluon propagator, $D_{\mu\nu}(k)$, the dressed quark-gluon vertex, $\Gamma_\nu^a(k, p)$, and λ^a are the usual SU(3) colour matrices. The solution to the gap equation (9) reads

$$\begin{aligned} S(p) &= -i\gamma \cdot p \sigma_V(p^2) + \sigma_S(p^2) \\ &= [i\gamma \cdot p A(p^2) + B(p^2)]^{-1}. \end{aligned} \quad (11)$$

The renormalisation constants for the quark-gluon vertex, $Z_1(\zeta, \Lambda^2)$, and quark-wave function, $Z_2(\zeta, \Lambda^2)$, depend on the renormalisation point, ζ , the regularisation scale, Λ , and the gauge parameter, whereas the mass function $M(p^2) = B(p^2)/A(p^2)$ is independent of ζ . Since QCD is asymptotically free, it is useful to impose at large spacelike ζ^2 the renormalisation condition,

$$S^{-1}(p)|_{p^2=\zeta^2} = i\gamma \cdot p + m(\zeta^2), \quad (12)$$

where $m(\zeta^2)$ is the renormalised running quark mass, so that for $\zeta^2 \gg \Lambda_{\text{QCD}}^2$ quantitative matching with pQCD results can be made.

Infrared dressing of light quarks has *profound* consequences for hadron phenomenology [29]: the quark-wave function renormalisation, $Z(p^2) = 1/A(p^2)$, is suppressed whereas the dressed quark-mass function, $M(p^2) = B(p^2)/A(p^2)$, is enhanced in the infrared which expresses dynamical chiral symmetry breaking (DCSB) and is crucial to the emergence of a constituent quark mass scale. Both, numerical solutions of the quark DSE and simulations of lattice-regularised QCD [30], predict this behaviour of $M(p^2)$ and pointwise agreement between DSE and lattice results has been explored in Refs. [31,32]. Studies that do not implement light-quark dressing run into artefacts caused by rather *large* light-quark masses [17,19] to emulate confinement. This is because unphysical thresholds in transition amplitudes can only be overcome with the prescription that $m_H < m_{q_1} + m_{q_2}$, which poses problems for a description of light vector mesons (ρ, K^*), heavy flavoured vector mesons (D^*, B^*) and for P -wave and excited charmonium states.

Whereas the impact of gluon dressing is striking for light quarks, its effect on the heavy quarks is barely notable. This can be appreciated, for instance, in Fig. 1 of Ref. [28]: for light quarks,

mass can be generated from nothing, *i.e.*, the Higgs mechanism is irrelevant to their acquiring of a constituent-like mass.

Bethe-Salpeter amplitudes: The BSA can be determined reliably by solving the Bethe-Salpeter equation (BSE) in a truncation scheme consistent with that employed in the gap equation (9). Consider the inhomogeneous BSE for the axialvector vertex $\Gamma_{5\mu}^{fg}$ in which pseudoscalar and axialvector mesons appear as poles:

$$\begin{aligned} \Gamma_{5\mu}^{fg}(k; P) &= Z_2 \gamma_5 \gamma_\mu - g^2 \int_q^\Lambda D^{\alpha\beta}(k-q) \frac{\lambda^a}{2} \gamma_\alpha \\ &\times S_f(q_+) \Gamma_{5\mu}^{fg}(q; P) S_g(q_-) \frac{\lambda^a}{2} \Gamma_\beta^g(q_-, k_-) \\ &+ g^2 \int_q^\Lambda D^{\alpha\beta}(k-q) \frac{\lambda^a}{2} \gamma_\alpha S_f(q_+) \frac{\lambda^a}{2} \Lambda_{5\mu\beta}^{fg}(k, q; P). \end{aligned} \quad (13)$$

In Eq. (13), P is the total meson momentum, $q_\pm = q \pm P/2$, $k_\pm = k \pm P/2$, $\Lambda_{5\mu\beta}^{fg}$ is a 4-point Schwinger function entirely defined via the quark self energy [33] and f, g denote the flavour indices of a light-light or heavy-light bound state. The solutions of the vertex $\Gamma_{5\mu}^{fg}$ must satisfy the axial-vector Ward-Takahashi identity,

$$\begin{aligned} P^\mu \Gamma_{5\mu}^{fg}(k; P) &= S_f^{-1}(k_+) i\gamma_5 + i\gamma_5 S_g^{-1}(k_-) \\ &- i[m_f(\zeta) + m_g(\zeta)] \Gamma_5^{fg}(k; P), \end{aligned} \quad (14)$$

where Γ_5^{fg} solves the pseudoscalar analogue to Eq. (13). A systematic, symmetry-preserving truncation of the DSE and BSE is given by the Rainbow ladder [34] which is their leading-order term with the dressed quark-gluon vertex, Γ_μ^f , replaced by γ_μ . It can be shown that $\Lambda_{5\mu\beta}^{fg} \equiv 0$ in this approximation.

The above Ward-Takahashi identity possesses another remarkable property; the set of quark-level Goldberger-Treiman relations that follow from it reveal the full structure of the light-meson BSA [35]. In particular, it enables one to relate the leading covariant, $\varepsilon_5(k; P)$, of the light pseudoscalar BSA with the scalar part, $B(p^2)$, of the dressed-quark propagator (11) in the chiral limit. This motivates an effective parametrisation of the light mesons ($M = \pi, K$),

$$\Gamma_M(k; P) = i\gamma_5 \varepsilon_M(k^2) = i\gamma_5 B_M(k^2)/\hat{f}_M, \quad (15)$$

$\hat{f}_M = f_M/\sqrt{2}$, which has been capitalised on in transition form factor calculations [20,27,28].

Simultaneous solutions of the quarks's DSE and the heavy meson's BSA with renormalisation-group improved ladder truncations, obtained for the kaon [35], prove to be difficult. The truncations do not yield the Dirac equation when *one* of the quark masses is large. A recent attempt to calculate BSA for D and B mesons [36] reproduces well the respective masses but underestimates experimental leptonic decay constants by 30 – 50%. With a consistent derivation of the heavy meson BSA pending, simple one-covariant forms for $\Gamma_H(k; P)$ are currently employed in Eq. (6), which reproduce leptonic decay constants in a simultaneous calculation.

6. Hadronic decays

The decay $D^* \rightarrow D\pi$ can be used to extract the strong coupling $\hat{g}$ between heavy vector and pseudoscalar mesons to a low-momentum pion in the heavy meson chiral lagrangian [37]. One considers the matrix element,

$$\langle D(p_2)\pi(q)|D^*(p_1, \lambda)\rangle = g_{D^*D\pi} \epsilon_{\lambda} \cdot q, \quad (16)$$

where the coupling, $g_{D^*D\pi} = 17.9 \pm 0.3 \pm 1.9$, is experimentally known [38] and related to $\hat{g}$. Similarly, one may also extract $\hat{g}$ from the unphysical decay $B^* \rightarrow B\pi$ in the chiral limit and where m_b/Λ_{QCD} corrections are better controlled.

The coupling $g_{D^*D\pi}$ is related to a heavy-to-heavy transition form factor via the LSZ reduction of the pion and the use of PCAC, $\pi(x) = \partial^\mu A_\mu(x)/(f_\pi m_\pi^2)$, which leads to:

$$\begin{aligned} \langle D(p_2)\pi(q)|D^*(p_1)\rangle &= q^\mu \frac{(m_\pi^2 - q^2)}{f_\pi m_\pi^2} \\ &\times \int d^4x e^{iq \cdot x} \langle D(p_2)|A_\mu(x)|D^*(p_1)\rangle. \end{aligned} \quad (17)$$

Hence, the matrix element in Eq. (16) has been reduced to the Fourier transform of a transition matrix element between the D^* and D mesons in the chiral limit with the axial QCD current $A_\mu(x) = \bar{q}^a \gamma_\mu \gamma_5 q^b$. Results from this reduction procedure have been obtained on the lattice [39, 40] and most recently in a simulation with $n_f = 2$ [41] which yields $g_{D^*D\pi} = 20 \pm 2$.

This form factor can also be calculated straightforwardly without reduction of the pion employing Eq. (6) with the dressed quark propagators in Eq. (11) and substituting the pion's BA (15) for Γ_L . In a reassessment and improvement of a calculation of $g_{D^*D\pi}$ within a Dyson-Schwinger model [28], we obtain $g_{D^*D\pi} = 21$ [42] in agreement with the lattice result [41] and about 16% larger than the experimental value.

7. Conclusive remarks

We have stressed the importance of hadronic effects in decays of heavy-flavoured mesons and portrayed the various theoretical *Ansätze* for the heavy-to-light transition form factors. In short, the main obstacle to their *precise* calculation, which veraciously reproduces the infrared features of QCD, are the uncertainties of the light-cone DA in the case of LCSR and the lack of model-independent wave functions in relativistic quark model calculations. We have argued that the running quark mass of the DSE quark propagators is crucial to include confinement and DCSB effects in the transition amplitudes; an unfinished task are consistent solutions of the BSE for the D and B mesons within the DSE formalism, which will reduce model dependence.

Acknowledgments

Based on the talk given at Light Cone 2009: Relativistic Hadronic and Particle Physics, 8–13 July 2009, São José dos Campos, São Paulo, Brazil. B. E. thanks the organisers at the Instituto Tecnológico de Aeronáutica for the welcoming atmosphere and in particular Tobias Frederico and João Pacheco de Melo for their hospitality. Several stimulating discussions with Arlene Aguilar, Adriano Natale, Fernando Navarra, Marina Nielsen, Joannis Papavassiliou and Lauro Tomio were greatly appreciated. This work was supported by the Department of Energy, Office of Nuclear Physics, No. DE-AC02-06CH11357.

REFERENCES

1. C. Bebek *et al.* [CLEO Collaboration], Phys. Rev. Lett. **46**, 84 (1981); K. Chadwick *et al.* [CLEO Collaboration], *ibid.* **46**, 88 (1981).

2. G. Buchalla, A. J. Buras and M. E. Lautenbacher, *Rev. Mod. Phys.* **68**, 1125 (1996).
3. M. Beneke, G. Buchalla, M. Neubert and C. T. Sachrajda, *Phys. Rev. Lett.* **83**, 1914 (1999); *id.*, *Nucl. Phys. B* **591**, 313 (2000).
4. Y. Y. Keum, H. N. Li and A. I. Sanda, *Phys. Rev. D* **63**, 054008 (2001).
5. C. W. Bauer *et al.*, *Phys. Rev. D* **63**, 114020 (2001); C. W. Bauer, D. Pirjol and I. W. Stewart, *ibid.* **65**, 054022 (2002); C. W. Bauer *et al.*, *ibid.* **66**, 014017 (2002).
6. C. W. Bauer *et al.*, *Phys. Rev. D* **70**, 054015 (2004).
7. M. Beneke and S. Jäger, *Nucl. Phys. B* **751**, 160 (2006); *id.*, *Nucl. Phys. B* **768**, 51 (2007).
8. G. P. Lepage and S. J. Brodsky, *Phys. Rev. D* **22**, 2157 (1980).
9. B. El-Bennich *et al.*, *Phys. Rev. D* **79**, 094005 (2009); B. El-Bennich *et al.*, *Phys. Rev. D* **74**, 114009 (2006).
10. M. Artuso *et al.*, *Eur. Phys. J. C* **57**, 309 (2008).
11. D. Bećirević and A. B. Kaidalov, *Phys. Lett. B* **478**, 417 (2000).
12. D. Ebert, R. N. Faustov and V. O. Galkin, *Phys. Rev. D* **75**, 074008 (2007).
13. M. A. Ivanov, J. G. Körner and P. Santorelli, *Phys. Rev. D* **63**, 074010 (2001); *id.* **71**, 094006 (2005); **75**, 019901(E) (2007).
14. A. Faessler *et al.*, *Eur. Phys. J. direct C* **4**, 18 (2002).
15. M. A. Ivanov, J. G. Körner and O. N. Pakhomova, *Phys. Lett. B* **555**, 189 (2003).
16. D. Melikhov, N. Nikitin and S. Simula, *Phys. Rev. D* **57**, 6814 (1998); D. Melikhov, *Eur. Phys. J. direct C* **2**, 1 (2002).
17. B. El-Bennich, O. Leitner, J. P. Dedonder and B. Loiseau, *Phys. Rev. D* **79**, 076004 (2009).
18. C. D. Lu, W. Wang and Z. T. Wei, *Phys. Rev. D* **76**, 014013 (2007).
19. B. El-Bennich *et al.*, *Braz. J. Phys.* **38**, 465 (2008).
20. M. A. Ivanov *et al.*, *Phys. Rev. D* **76**, 034018 (2007).
21. F. De Fazio, T. Feldmann and T. Hurth, *Nucl. Phys. B* **733**, 1 (2006) [Erratum-*ibid.* **B 800**, 405 (2008)].
22. A. Khodjamirian, T. Mannel and N. Offen, *Phys. Lett. B* **620**, 52 (2005); *id.*, *Phys. Rev. D* **75**, 054013 (2007).
23. P. Ball and R. Zwicky, *JHEP* **0110**, 019 (2001); *id.*, *Phys. Rev. D* **71**, 014015 (2005).
24. P. A. Boyle *et al.*, [UKQCD Collab.], *Phys. Lett. B* **641**, 67 (2006).
25. C. D. Roberts and A. G. Williams, *Prog. Part. Nucl. Phys.* **33**, 477 (1994).
26. C. D. Roberts, M. S. Bhagwat, A. Holl and S. V. Wright, *Eur. Phys. J. ST* **140**, 53 (2007).
27. M. A. Ivanov *et al.*, *Phys. Rev. C* **57**, 1991 (1998).
28. M. A. Ivanov, Yu. L. Kalinovsky and C. D. Roberts, *Phys. Rev. D* **60**, 034018 (1999).
29. C. D. Roberts, *Nucl. Phys. A* **605**, 475 (1996); C. D. Roberts and S. M. Schmidt, *Prog. Part. Nucl. Phys.* **45**, S1 (2000); R. Alkofer and L. von Smekal, *Phys. Rept.* **353**, 281 (2001).
30. J. B. Zhang *et al.*, *Phys. Rev. D* **71**, 014501 (2005).
31. M. S. Bhagwat, M. A. Pichowsky, C. D. Roberts and P. C. Tandy, *Phys. Rev. C* **68**, 015203 (2003); M. S. Bhagwat and P. C. Tandy, *Phys. Rev. D* **70**, 094039 (2004).
32. R. Alkofer, W. Detmold, C. S. Fischer and P. Maris, *Phys. Rev. D* **70**, 014014 (2004).
33. L. Chang and C. D. Roberts, *Phys. Rev. Lett.* **103**, 081601 (2009).
34. H. J. Munczek, *Phys. Rev. D* **52**, 4736 (1995); A. Bender, C. D. Roberts and L. Von Smekal, *Phys. Lett. B* **380**, 7 (1996).
35. P. Maris, C. D. Roberts and P. C. Tandy, *Phys. Lett. B* **420**, 267 (1998); P. Maris and C. D. Roberts, *Phys. Rev. C* **56**, 3369 (1997).
36. T. Nguyen, N. A. Souchlas and P. C. Tandy, *AIP Conf. Proc.* **1116**, 327 (2009).
37. M. B. Wise, *Phys. Rev. D* **45**, 2188 (1992).
38. A. Anastassov *et al.* [CLEO Collaboration], *Phys. Rev. D* **65**, 032003 (2002).
39. G. M. de Divitiis *et al.* [UKQCD Collaboration], *JHEP* **9810**, 010 (1998).
40. A. Abada *et al.*, *Phys. Rev. D* **66**, 074504 (2002).
41. D. Bećirević and B. Haas, arXiv:0903.2407 [hep-lat]; D. Bećirević, B. Blossier, E. Chang and B. Haas, *Phys. Lett. B* **679**, 231 (2009).
42. B. El-Bennich, M. A. Ivanov and C. D. Roberts, in preparation.